



## 青蔥病蟲害防治藥劑

| 病蟲害別 | 藥劑名稱   | 稀釋倍數  | 安全採收期(天) | 作用機制代號       |
|------|--|-------|----------|--------------|
| 白絹病  | 50%撲滅寧水懸劑                                      | 2,000 | 14       | 2, E3        |
|      | 50%撲滅寧可溼性粉劑                                    | 2,000 | 14       | 2, E3        |
|      | 50%大克爛可溼性粉劑                                    | 2,000 | 12       | 14, F3       |
|      | 15%福多寧乳劑                                       | 900   | 18       | 7, C2        |
|      | 20%福多寧水懸劑                                      | 1,200 | 18       | 7, C2        |
|      | 50%福多寧可溼性粉劑                                    | 3,000 | 18       | 7, C2        |
| 灰黴病  | 42.4%白克列水懸劑                                    | 1,200 | 14       | 7, C2        |
|      | 50%白克列水分散性粒劑                                   | 1,500 | 14       | 7, C2        |
|      | 1x10 <sup>9</sup> CFU/g 貝萊斯芽孢桿菌 BF 可溼性粉劑       | 600   | -        | BM02         |
|      | >1x10 <sup>9</sup> CFU/mL 液化澱粉芽孢桿菌 Ba-BPD1 水懸劑 | 300   | -        | BM02         |
|      | >1x10 <sup>8</sup> CFU/mL 液化澱粉芽孢桿菌 CL3 水懸劑     | 400   | -        | BM02         |
| 立枯病  | 25%賓克隆可溼性粉劑                                    | 1,000 | 15       | 20, B4       |
|      | 23.2%賓克隆水懸劑                                    | 1,000 | 15       | 20, B4       |
|      | 25%賓克隆水懸劑                                      | 1,000 | 15       | 20, B4       |
| 炭疽病  | 27.3%三氟得克利水懸劑                                  | 2,500 | 21       | 11, C3+3, G1 |
|      | 24.9%待克利乳劑                                     | 3,000 | 21       | 3, G1        |
|      | 24.9%待克利水懸劑                                    | 3,000 | 21       | 3, G1        |
|      | 10%待克利水分散性粒劑                                   | 1,200 | 21       | 3, G1        |
|      | 250g/L 待克利乳劑                                   | 3,000 | 21       | 3, G1        |
|      | 75%四氯異苯腈可溼性粉劑                                  | 600   | 7        | M5, mc5      |
|      | 75%四氯異苯腈水分散性粒劑                                 | 600   | 7        | M5, mc5      |
|      | 40%四氯異苯腈水懸劑                                    | 320   | 7        | M5, mc5      |
|      | 75%四氯異苯腈可溼性粉                                   | 600   | 7        | M5, mc5      |



| 病蟲害別   | 藥劑名稱   | 稀釋倍數  | 安全採收期(天) | 作用機制代號         |
|--------|--|-------|----------|----------------|
|        | 劑水溶性袋裝   |       |          |                |
|        | 23.6%百克敏乳劑                                     | 3,000 | 9        | 11, C3         |
|        | 23.6%百克敏水懸劑                                    | 3,000 | 9        | 11, C3         |
|        | 1x10 <sup>9</sup> CFU/g 貝萊斯芽孢桿菌 BF 可濕性粉劑       | 600   | -        | BM02           |
| 疫病     | 27.12%三元硫酸銅水懸劑                                 | 500   | -        | M1, mc1        |
|        | 81.3%嘉賜銅可溼性粉劑                                  | 800   | 6        | 24, D3+M1, mc1 |
|        | 9.4%賽座滅水懸劑                                     | 2,000 | 9        | 21, C4         |
|        | 35%滅達樂種子處理水分散性粉劑                               | -     | -        | 4, A1          |
|        | 17.5%滅達樂種子處理液劑                                 | -     | -        | 4, A1          |
|        | 17.5%滅達樂溶液                                     | -     | -        | 4, A1          |
|        | 35%滅達樂水懸劑                                      | -     | -        | 4, A1          |
|        | 35%滅達樂可溼性粉劑                                    | -     | -        | 4, A1          |
|        | 2.5%右滅達樂粒劑                                     | -     | -        | 4, A1          |
|        | 50%亞托敏水分散性粒劑                                   | 2,000 | 10       | 11, C3         |
|        | 10%亞托敏水懸劑                                      | 400   | 10       | 11, C3         |
|        | 23%亞托敏水懸劑                                      | 1000  | 10       | 11, C3         |
|        | 250g/L 亞托敏水懸劑                                  | 1000  | 10       | 11, C3         |
|        | 50%亞托敏水分散性粒劑                                   | 2000  | 10       | 11, C3         |
|        | 60.8%氟比拔克水懸劑                                   | 1000  | 21       | 28, F4+43,B5   |
|        | 250g/L 曼普胺水懸劑                                  | 2500  | 3        | 40, H5         |
|        | 1X10 <sup>7</sup> CFU/g 蓋棘木黴菌 ICC080/012 可濕性粉劑 | 2000  | -        | BM02           |
| 細菌性軟腐病 | 27.12%三元硫酸銅水懸劑                                 | 500   | -        | M1, mc1        |
|        | 81.3%嘉賜銅可溼性粉劑                                  | 800   | 6        | 24, D3+M1, mc1 |
|        | 14%嘉賜克枯爛可溼性粉                                   | 1,000 | 6        | 24, D3+34, un  |



| 病蟲害別 | 藥劑名稱                          | 稀釋倍數  | 安全採收期(天) | 作用機制代號         |
|------|-------------------------------|-------|----------|----------------|
|      | 劑                             |       |          |                |
|      | 68.8%多保鏈黴素可溼性粉劑               | 1,000 | 12       | 1, B1+25, D4   |
|      | 12.5%鏈黴素溶液                    | 1,000 | -        | 25, D4         |
| 紫斑病  | 50%依普同可溼性粉劑                   | 1,000 | 6        | 2, E3          |
|      | 23.7%依普同水懸劑                   | 1,000 | 15       | 2, E3          |
|      | 23%亞托敏水懸劑                     | 3,000 | 10       | 11, C3         |
|      | 250 g/L 亞托敏水懸劑                | 3000  | 10       | 11, C3         |
|      | 50%亞托敏水分散性粒劑                  | 6,000 | 10       | 11, C3         |
|      | 10%亞托敏水懸劑                     | 1,200 | 10       | 11, C3         |
|      | 38%白列克敏水分散性粒劑                 | 3,000 | 10       | 11, C3+7, C2   |
|      | 81.3%嘉賜銅可溼性粉劑                 | 800   | 6        | 24, D3+M1, mc1 |
|      | 45%甲鋅保粒素可溼性粉劑                 | 500   | 10       | 19, H4+M3, mc3 |
|      | 44.2%克收欣水懸劑                   | 2,500 | 10       | 11, C3         |
|      | 50%克收欣水分散性粒劑                  | 2,800 | 10       | 11, C3         |
|      | 40%四氯異苯腈水懸劑                   | 320   | 7        | M5, mc5        |
|      | 75%四氯異苯腈水分散性粒劑                | 700   | 7        | M5, mc5        |
|      | 75%四氯異苯腈可溼性粉劑                 | 1,500 | 7        | M5, mc5        |
|      | 75%四氯異苯腈可溼性粉劑水溶性袋裝            | 700   | 7        | M5, mc5        |
|      | 21.2%依滅列乳劑                    | 1,500 | 6        | 3, G1          |
|      | 10%保粒黴素(甲)可溼性粉劑               | 1,000 | -        | 19, H4         |
|      | 50%保粒黴素(甲)水溶性粒劑               | 5,000 | -        | 19, H4         |
|      | ≥5*10 <sup>9</sup> cfu/g 液化澱粉 | 600   | -        | BM02           |



| 病蟲害別            | 藥劑名稱             | 稀釋倍數        | 安全採收期(天) | 作用機制代號        |
|-----------------|------------------|-------------|----------|---------------|
|                 | 芽孢桿菌 YCMA1 可溼性粉劑 |             |          |               |
| 銹病              | 23%三泰隆水分散性乳劑     | 2,000       | 15       | 3, G1         |
|                 | 23%三泰隆乳劑         | 2,000       | 15       | 3, G1         |
|                 | 43.7%三氟敏水懸劑      | 3,500       | 12       | 11, C3        |
|                 | 50%三氟敏水分散性粒劑     | 4,000       | 12       | 11, C3        |
|                 | 500g/L 三氟敏水懸劑    | 4,000       | 12       | 11, C3        |
|                 | 84.2%三得芬乳劑       | 2,000       | 6        | 5, G2         |
|                 | 25%普克利乳劑         | 1,000       | 14       | 3, G1         |
|                 | 250g/L 普克利乳劑     | 1,000       | 14       | 3, G1         |
|                 | 5%菲克利水懸劑         | 1,500       | 9        | 3, G1         |
|                 | 23%菲克利水懸劑        | 6,900       | 9        | 3, G1         |
|                 | 10%菲克利水懸劑        | 3,000       | 9        | 3, G1         |
|                 | 10%菲克利乳劑         | 3,000       | 9        | 3, G1         |
|                 | 23.6%百克敏乳劑       | 2,500       | 9        | 11, C3        |
|                 | 23.6%百克敏水懸劑      | 2,500       | 9        | 11, C3        |
|                 | 28%亞托環克座水懸劑      | 3,000       | 10       | 11, C3+3, G1  |
|                 | 露菌病              | 80%錳乃浦可溼性粉劑 | 400      | 7             |
| 37%錳乃浦水懸劑       |                  | 200         | 7        | M3, mc3       |
| 71.6%銅右滅達樂可溼性粉劑 |                  | 600         | 3        | 4, A1+M1, mc1 |
| 23%亞托敏水懸劑       |                  | 2,000       | 10       | 11, C3        |
| 250g/L 亞托敏水懸劑   |                  | 2,000       | 10       | 11, C3        |
| 50%亞托敏水分散性粒劑    |                  | 4,500       | 10       | 11, C3        |
| 10%亞托敏水懸劑       |                  | 900         | 10       | 11, C3        |
| 500g/L 達滅芬水懸劑   |                  | 3,000       | 14       | 40, H5        |
| 50%達滅芬水懸劑       |                  | 3,000       | 14       | 40, H5        |
| 50%達滅芬可溼性粉劑     |                  | 3,000       | 14       | 40, H5        |
| 76.5%銅滅達樂可溼性粉劑  |                  | 600         | 3        | 4, A1+M1, mc1 |



| 病蟲害別 | 藥劑名稱   | 稀釋倍數  | 安全採收期(天) | 作用機制代號       |
|------|--|-------|----------|--------------|
|      | 48%松香酯銅乳劑                                    | 1000  | -        | -            |
|      | 60.8%氟比拔克水懸劑                                 | 1500  | 21       | 28, F4+43,B5 |
|      | >1x10 <sup>9</sup> CFU/g 液化澱粉芽孢桿菌 QST713 水懸劑 | 400   | -        | BM02         |
| 夜蛾類  | 300g/L(30%w/v)賽速安勃水懸劑                        | 4,000 | 9        | 28+4A        |
|      | 150g/L 賽洛安勃膠囊水懸混劑                            | 4,000 | 9        | 28+3A        |
|      | 70%得芬諾可溼性粉劑                                  | 7,000 | 9        | 18           |
|      | 19.7%得芬諾水懸劑                                  | 2,000 | 9        | 18           |
|      | 5%因滅汀乳劑                                      | 5,000 | 9        | 6            |
|      | 19.2g/L(1.92%w/v)因滅汀乳劑                       | 1,500 | 9        | 6            |
|      | 2.15%因滅汀水基乳劑                                 | 1,500 | 9        | 6            |
|      | 2.15%因滅汀乳劑                                   | 1,500 | 9        | 6            |
|      | 2.15%因滅汀水懸劑                                  | 1,500 | 9        | 6            |
|      | 5%因滅汀水溶性粒劑                                   | 5,000 | 9        | 6            |
|      | 14.5%因得克水懸劑                                  | 2,000 | 9        | 22A          |
|      | 15.8%因得克乳劑                                   | 2,000 | 9        | 22A          |
|      | 5%可芬諾水懸劑                                     | 1,000 | 18       | 18           |
|      | 2.5%畢芬寧水懸劑                                   | 1,000 | 14       | 3A           |
|      | 2.8%畢芬寧乳劑                                    | 1,000 | 14       | 3A           |
|      | 18.4%剋安勃水懸劑                                  | 2,500 | 9        | 28           |
|      | 35%剋安勃水分散性粒劑                                 | 5,000 | 9        | 28           |
|      | 54%庫斯蘇力菌 ABTS-351 水分散性粒劑                     | 1500  | -        | 11A          |
|      | 48.1%(35,000DBMU/MG) 鮎澤蘇力菌 ABTS-1857 水分散性粒劑  | 1,000 | -        | 11A          |
|      | 9.6%氟芬隆水分散性乳劑                                | 3,000 | 12       | 15           |



| 病蟲害別 | 藥劑名稱                    | 稀釋倍數  | 安全採收期(天) | 作用機制代號 |
|------|-------------------------|-------|----------|--------|
|      | 100g/L(10%w/v)氟芬隆水分散性乳劑 | 3,000 | 12       | 15     |
|      | 4.4%祿芬隆乳劑               | 1,500 | 10       | 15     |
|      | 50g/L(5%w/v)祿芬隆乳劑       | 1,500 | 10       | 15     |
|      | 20%氟大滅水分散性粒劑            | 3,000 | 21       | 28     |
|      | 5%賽滅寧乳劑                 | 1,500 | 6        | 3A     |
|      | 5%賽滅寧微乳劑                | 1,500 | 6        | 3A     |
|      | 5%賽滅寧水基乳劑               | 1,500 | 6        | 3A     |
|      | 5%賽滅寧可溼性粉劑              | 1,500 | 6        | 3A     |
|      | 10%賽滅寧可溼性粉劑             | 3,000 | 6        | 3A     |
|      | 50%培丹水溶性粉劑              | 1,000 | 10       | 14     |
|      | 40%納乃得水溶性粉劑-水溶性袋裝       | 1,500 | 4        | 1A     |
|      | 40%納乃得水溶性粉劑             | 1,500 | 4        | 1A     |
|      | 25%納乃得水溶性粉劑             | 900   | 4        | 1A     |
|      | 40%納乃得水溶性粒劑             | 1,500 | 4        | 1A     |
|      | 10%克凡派水懸劑               | 1,000 | 9        | 13     |
|      | 58%乃力松乳劑                | 1000  | 4        | 1B     |
|      | 5%克福隆乳劑                 | 2000  | 12       | 15     |
|      | 20%芬化利乳劑                | 3000  | 14       | 3A     |
|      | 5%芬化利乳劑                 | 750   | 14       | 3A     |
|      | 20%芬化利水基乳劑              | 3000  | 14       | 3A     |
|      | 20%芬化利可濕性粉劑             | 3000  | 14       | 3A     |
|      | 10%芬化利乳劑                | 1500  | 14       | 3A     |
|      | 6%苦參鹼溶液                 | 1000  | -        | -      |
|      | 40%益滅賽寧可濕性粉劑            | 2500  | 6        | 1B+3A  |
|      | 13.5%納得亞滅寧乳劑            | 1000  | 6        | 1A+3A  |
|      | 50%馬拉松乳劑                | 500   | 4        | 1B     |
|      | 50%馬拉松水基乳劑              | 500   | 4        | 1B     |
|      | 25%馬拉松可濕性粉劑             | 250   | 4        | 1B     |





| 病蟲害別 | 藥劑名稱                | 稀釋倍數                                   | 安全採收期(天) | 作用機制代號 |
|------|---------------------|--|----------|--------|
|      | 50%馬拉松可濕性粉劑         | 500                                    | 4        | 1B     |
|      | 34%硫敵克水懸劑           | 1000                                   | 6        | 1A     |
|      | 75%硫敵克可濕性粉劑         | 2500                                   | 6        | 1A     |
|      | 2.8%第滅寧乳劑           | 1500                                   | 6        | 3A     |
|      | 25g/L 第滅寧乳劑         | 1500                                   | 6        | 3A     |
|      | 2.8%第滅寧水基乳劑         | 1500                                   | 6        | 3A     |
|      | 2.4%第滅寧水懸劑          | 1500                                   | 6        | 3A     |
|      | 2.46%賽洛寧膠囊懸著劑       | 1000                                   | 10       | 3A     |
|      | 2.8%賽洛寧乳劑           | 2000                                   | 9        | 3A     |
|      | 2.5%賽洛寧微乳劑          | 2000                                   | 9        | 3A     |
|      | 2.8%賽洛寧水懸劑          | 1000                                   | 10       | 3A     |
|      | 5%賽洛寧水分散性粒劑         | 2000                                   | 10       | 3A     |
|      | 1%賽洛寧可濕性粉劑          | 400                                    | 10       | 3A     |
|      | 甜菜夜蛾                | 2x10 <sup>9</sup> OBs/ml 甜菜夜蛾核多角體病毒水懸劑 | 3,000    | -      |
| 蔥潛蠅  | 10%福瑞松粒劑            | 種植前使用                                  | -        | 1B     |
|      | 50%益滅松可溼性粉劑         | 1,000                                  | 7        | 1B     |
|      | 8.9%賽滅淨溶液           | 800                                    | 7        | 17     |
|      | 75%賽滅淨可溼性粉劑         | 5000                                   | 7        | 17     |
| 潛蠅類  | 8.9%賽滅淨溶液           | 800                                    | 7        | 17     |
|      | 75%賽滅淨可溼性粉劑         | 5,000                                  | 7        | 17     |
| 蚜蟲類  | 31.6%護賽寧乳劑          | 8,500                                  | 15       | 3A     |
|      | 20%免扶克乳劑            | 500                                    | 12       | 1A     |
|      | 2.4%第滅寧水懸劑          | 1,000                                  | 6        | 3A     |
|      | 2.8%第滅寧乳劑           | 1,000                                  | 6        | 3A     |
|      | 2.8%第滅寧水基乳劑         | 1,000                                  | 6        | 3A     |
|      | 25g/L(2.5%w/v)第滅寧乳劑 | 1000                                   | 6        | 3A     |
|      | 50%培丹水溶性粉劑          | 1,000                                  | 10       | 14     |



| 病蟲害別 | 藥劑名稱              | 稀釋倍數  | 安全採收期(天) | 作用機制代號 |
|------|-------------------|-------|----------|--------|
|      | 58%乃力松乳劑          | 1,000 | 4        | 1B     |
|      | 1%密滅汀乳劑           | 1500  | 6        | 6      |
|      | 50%賽達松乳劑          | 1000  | 15       | 1B     |
|      | 50%賽達松水基乳劑        | 1000  | 15       | 1B     |
|      | 28.8%益達胺溶液        | 12000 | 9        | 4A     |
|      | 18.2%益達胺水懸劑       | 8000  | 9        | 4A     |
|      | 9.6%益達胺溶液         | 4000  | 9        | 4A     |
|      | 9.6%益達胺水懸劑        | 4000  | 9        | 4A     |
|      | 25%納乃得水溶性粉劑       | 800   | 4        | 1A     |
|      | 40%納乃得水溶性粒劑       | 1500  | 4        | 1A     |
|      | 40%納乃得水溶性粉劑-水溶性袋裝 | 1500  | 4        | 1A     |
|      | 40%納乃得水溶性粉劑       | 1,500 | 4        | 1A     |
|      | 25%免扶克可溼性粉劑       | 700   | 12       | 1A     |
|      | 40%免扶克水懸劑         | 1,000 | 12       | 1A     |
|      | 25%賽速安水溶性粒劑       | 5,000 | 9        | 4A     |
|      | 10%賽速安水溶性粒劑       | 2,000 | 9        | 4A     |
|      | 21.8%速殺氟水懸劑       | 6,000 | 6        | 4C     |
|      | 20%達特南水溶性粒劑       | 2,000 | 7        | 4A     |
| 薊馬類  | 2.8%畢芬寧乳劑         | 1,000 | 14       | 3A     |
|      | 2.5%畢芬寧水懸劑        | 1,000 | 14       | 3A     |
|      | 10%克凡派水懸劑         | 1,000 | 9        | 13     |
|      | 5%賽滅寧乳劑           | 1,500 | 6        | 3A     |
|      | 5%賽滅寧微乳劑          | 1,500 | 6        | 3A     |
|      | 5%賽滅寧水基乳劑         | 1,500 | 6        | 3A     |
|      | 5%賽滅寧可溼性粉劑        | 1,500 | 6        | 3A     |
|      | 10%賽滅寧可溼性粉劑       | 3,000 | 6        | 3A     |
|      | 28.8%益達胺溶液        | 6,000 | 9        | 4A     |





| 病蟲害別  | 藥劑名稱                  | 稀釋倍數  | 安全採收期(天) | 作用機制代號 |
|-------|-----------------------|-------|----------|--------|
|       | 18.2%益達胺水懸劑           | 4,000 | 9        | 4A     |
|       | 9.6%益達胺溶液             | 2,000 | 9        | 4A     |
|       | 9.6%益達胺水懸劑            | 2,000 | 9        | 4A     |
|       | 2.9%貝他 - 賽扶寧乳劑        | 1,500 | 9        | 3A     |
|       | 2.5%賽洛寧微乳劑            | 2,000 | 9        | 3A     |
|       | 2.46%賽洛寧膠囊懸著劑         | 2,000 | 9        | 3A     |
|       | 2.8%賽洛寧乳劑             | 2,000 | 9        | 3A     |
|       | 2.8%賽洛寧水懸劑            | 2,000 | 9        | 3A     |
|       | 5%賽洛寧水分散性粒劑           | 3,500 | 9        | 3A     |
|       | 1%賽洛寧可溼性粉劑            | 700   | 9        | 3A     |
|       | 25%福化利乳劑              | 3,000 | 7        | 3A     |
|       | 240g/L(24%w/v)福化利水基乳劑 | 3,000 | 7        | 3A     |
|       | 500g/L(50%w/v)佈飛松乳劑   | 800   | 15       | 1B     |
|       | 43%佈飛松乳劑              | 800   | 15       | 1B     |
|       | 5.87%賜諾特水懸劑           | 4,000 | 6        | 5      |
|       | 11.7%賜諾特水懸劑           | 8,000 | 6        | 5      |
|       | 11.6%賜諾殺水懸劑           | 4500  | 3        | 5      |
|       | 2.5%賜諾殺水懸劑            | 1000  | 3        | 5      |
|       | 80%賜諾殺可溼性粉劑           | 32000 | 3        | 5      |
| 葉部薊馬類 | 28.8%益達胺溶液            | 6,000 | 6        | 4A     |
|       | 18.2%益達胺水懸劑           | 4,000 | 6        | 4A     |
|       | 9.6%益達胺溶液             | 2,000 | 6        | 4A     |
|       | 9.6%益達胺水懸劑            | 2,000 | 6        | 4A     |
|       | 50%滅賜克可溼性粉劑           | 800   | 12       | 1A     |
|       | 2.5%賽洛寧微乳劑            | 2,000 | 9        | 3A     |
|       | 2.46%賽洛寧膠囊懸著劑         | 2,000 | 9        | 3A     |
|       | 2.8%賽洛寧乳劑             | 2,000 | 9        | 3A     |
|       | 2.8%賽洛寧水懸劑            | 2,000 | 9        | 3A     |



| 病蟲害別      | 藥劑名稱                  | 稀釋倍數         | 安全採收期(天)                          | 作用機制代號        |
|-----------|-----------------------|--------------|-----------------------------------|---------------|
|           | 5%賽洛寧水分散性粒劑           | 4,000        | 9                                 | 3A            |
|           | 1%賽洛寧可溼性粉劑            | 800          | 9                                 | 3A            |
|           | 25%福化利乳劑              | 3,000        | 7                                 | 3A            |
|           | 240g/L(24%w/v)福化利水基乳劑 | 3,000        | 7                                 | 3A            |
| 毒蛾類       | 300g/L(30%w/v)賽速安勃水懸劑 | 4,000        | 9                                 | 28+4A         |
| 根蟻類       | 43%佈飛松乳劑              | 1,000        | 15                                | 1B            |
|           | 500g/L(50%w/v)佈飛松乳劑   | 1,000        | 15                                | 1B            |
| 鱗翅目<br>害蟲 | 2.8%畢芬寧乳劑             | 1,000        | 14                                | 3A            |
|           | 2.5%畢芬寧水懸劑            | 1,000        | 14                                | 3A            |
|           | 75%硫敵克可溼性粉劑           | 2,500        | 6                                 | 1A            |
|           | 34%硫敵克水懸劑             | 1,000        | 6                                 | 1A            |
|           | 5%芬化利乳劑               | 750          | 14                                | 3A            |
|           | 20%芬化利可溼性粉劑           | 3,000        | 14                                | 3A            |
|           | 20%芬化利水基乳劑            | 3,000        | 14                                | 3A            |
|           | 20%芬化利乳劑              | 3,000        | 14                                | 3A            |
|           | 10%芬化利乳劑              | 1,500        | 14                                | 3A            |
|           | 10%克凡派水懸劑             | 1000         | 10                                | 13            |
|           | 4.4%祿芬隆乳劑             | 1500         | 10                                | 15            |
|           | 50g/L(5%w/v)祿芬隆乳劑     | 1500         | 10                                | 15            |
|           | 150 g/L (15% w/v)賽洛安勃 | 4000         | 9                                 | 3A+28         |
|           | 切根蟲                   | 3%加福松微粒劑     | 50 公斤/<br>公頃                      | 定植前 3 天<br>施藥 |
| 0.5%芬化利粒劑 |                       | 50 公斤/<br>公頃 | 定植前 3 天<br>及定植後 3<br>天、10 天施<br>藥 | 3A            |



| 病蟲害別 | 藥劑名稱    | 稀釋倍數         | 安全採收期(天)      | 作用機制代號 |
|------|---------|--------------|---------------|--------|
|      | 5%免扶克粒劑 | 30 公斤/<br>公頃 | 定植前 3 天<br>施藥 | 1A     |

1. 上述資料更新日期以主管機關公告為準。其它防治藥劑及相關資料可參考主管機關之公告或參閱「農藥資訊服務網」--登記管理---病蟲害防治
2. 臺南區農業改良場病蟲害服務專線 06-5912959

1.5 版 112.09.11 整理